

334

2828 + small birds seen
que dice matagomes ~~Presente~~
y tomé sucedáneo ~~Pago~~ concauac
este aves tienen el color de alirán
en que se diferencia de las otras
que tienen la parte inferior blanca
que son ~~Podargus~~ donalle
algunas veces ~~Pavo~~
que tienen la parte inferior blanca
que son ~~Podargus~~ ~~de~~ ~~color~~ ~~gris~~ ~~o~~ ~~negro~~
otras que son ~~color~~ ~~gris~~ ~~o~~ ~~negro~~
y tienen la parte inferior blanca.

and the other is the lateral or transverse fold. The latter is the more common and is produced by the folding of the skin over the underlying bone. It is produced by the action of the platysma muscle which is a broad, thin, triangular muscle situated on the surface of the neck and extending downwards and forwards to cover the lower part of the sternum and the upper part of the xiphoid process. It is supplied by the internal thoracic artery and the external thoracic artery. The platysma muscle is divided into two parts, the anterior and the posterior. The anterior part is the larger and is situated on the anterior aspect of the sternum and the upper part of the xiphoid process. The posterior part is smaller and is situated on the posterior aspect of the sternum and the upper part of the xiphoid process. The platysma muscle is supplied by the internal thoracic artery and the external thoracic artery.

